

# Interaction of GTP-binding proteins with calmodulin

Tomiko Asano, Nobuaki Ogasawara, Satoko Kitajima\* and Mamoru Sano\*

*Department of Biochemistry and \*Department of Morphology, Institute for Developmental Research, Aichi Prefectural Colony, Kasugai, Aichi 480-03, Japan*

Received 6 May 1986

Two GTP-binding proteins ( $G_i$  and  $G_o$ ), which were the substrates for islet-activating protein, pertussis toxin, were purified from bovine cerebral cortical membranes. Both  $G_i$  and  $G_o$  completely inhibited calmodulin-stimulated cyclic nucleotide phosphodiesterase activity. The same concentrations of these proteins, however, had no appreciable effect on the basal phosphodiesterase activity. The isolated  $G_{i\alpha}$  and  $\beta\gamma$  subunits of GTP-binding proteins were potent inhibitors of the calmodulin-stimulated phosphodiesterase activity, but  $G_{o\alpha}$  was very weak. Therefore, the  $\beta\gamma$  subunits were likely to be the major active molecules in the brain membranes. GTP-binding proteins were shown to bind directly to calmodulin in a  $Ca^{2+}$ -dependent manner by a gel permeation binding experiment.

*GTP-binding protein      Calmodulin      Cyclic nucleotide phosphodiesterase      Protein interaction*

## 1. INTRODUCTION

Recent studies have demonstrated high concentrations of GTP-binding proteins in brain membranes, and two major proteins have been purified [1,2]. One is the inhibitory GTP-binding protein of adenylate cyclase ( $G_i$ ), and the other is an analogous protein termed  $G_o$  by Sternweis and Robishaw [1]. Both  $\alpha$  subunits bind GTP and are ADP-ribosylated by islet-activating protein, pertussis toxin. Both  $\beta\gamma$  subunits from the two proteins seem to be identical [1]. These two proteins account for 1.5% of the membrane proteins [1], and  $G_o$  exists several-times more than  $G_i$  in brain membranes. Molecular mechanisms of the action of these GTP-binding proteins have not been clear except for the role of  $G_i$  in adenylate cyclase system [3].

Calmodulin is a  $Ca^{2+}$ -binding protein that modulates the action of numerous  $Ca^{2+}$ -dependent processes [4]. It is found in high concentrations in the central nervous system. Calmodulin exists in both cytoplasmic and particulate fractions of cells [5] and is associated with specific membranous components of neurons, including synaptic [6] and

vesicular membranes [7]. Functionally, calmodulin has been shown to activate cyclic nucleotide phosphodiesterase [8], adenylate cyclase [9], and  $Ca^{2+}$ ,  $Mg^{2+}$ -ATPase [6] in brain. It has also been shown to be involved in  $Ca^{2+}$ -dependent phosphorylation of synaptic membranes [7], in axonal transport [10], and in the release of neurotransmitters [7].

In addition to calmodulin-regulated enzymes, various proteins and peptides including cytoskeletal proteins [4], opioid peptides [11], hormones [12], insect venoms [13] and seminalplasmin [14] have been shown to bind calmodulin or/and to inhibit the activity of calmodulin in a  $Ca^{2+}$ -dependent manner. But the biological significance of the binding of these proteins or peptides to calmodulin has not been clarified.

This study shows that GTP-binding proteins inhibit the calmodulin-stimulated phosphodiesterase activity and that this inhibition is apparently due to a direct  $Ca^{2+}$ -dependent interaction of GTP-binding proteins with calmodulin.

## 2. MATERIALS AND METHODS

Calmodulin was purified from rat testis by the method of Yazawa et al. [15]. [ $^3\text{H}$ ]Calmodulin was prepared by acetylation of rat testis calmodulin using [ $^3\text{H}$ ]acetic anhydride (Amersham) [16].

Calmodulin-deficient phosphodiesterase was prepared from the soluble fraction of bovine cerebral cortex as described by Sharma and Wang [17]. Cyclic nucleotide phosphodiesterase activity was measured by the method described in [18].

$G_i$  and  $G_o$  were purified from bovine cerebral cortex according to Sternweis and Robishaw [1]. Each subunit was resolved according to the method described in [19,20]. An equal amount of  $\beta\gamma$  subunits was added to  $G_{i\alpha}$  and  $G_{o\alpha}$  preparations to form  $G_i$  and  $G_o$ , respectively. To determine GTP-binding protein, the binding of [ $^{35}\text{S}$ ]guanosine 5'-(3-*O*-thio)triphosphate ( $\text{GTP}\gamma\text{S}$ ) (New England Nuclear) was measured essentially according to Northup et al. [21].

The binding of GTP-binding proteins to calmodulin was examined by the gel permeation binding technique of Hummel and Dreyer [22]. The preparation of  $G_i$  or  $G_o$  was subjected to Sephadex G-50 column (0.7  $\times$  11 cm) equilibrated with [ $^3\text{H}$ ]calmodulin (10000 cpm/ml) in 10 mM Hepes (pH 8.0), 0.5 mM dithiothreitol (DTT), 0.05% Lubrol PX, and 0.5 mM  $\text{CaCl}_2$  or 0.5 mM EGTA, and the radioactivity of each fraction (105  $\mu\text{l}$ ) was counted.

## 3. RESULTS

Fig.1 shows the effects of GTP-binding proteins on cyclic nucleotide phosphodiesterase activity. Both  $G_i$  and  $G_o$  inhibited calmodulin-stimulated phosphodiesterase activity with  $\text{IC}_{50}$  values of 130 and 760 nM, respectively, but the same concentrations of these proteins had no significant effect on the basal phosphodiesterase activity. The effect of each subunit of  $G_i$  and  $G_o$  was also determined (fig.1).  $G_{i\alpha}$  and  $\beta\gamma$  subunits exerted potent inhibition on the calmodulin-stimulated activity but  $G_{o\alpha}$  subunit was a very weak inhibitor. Therefore, the inhibitory effect of  $G_o$  seemed to be due to the effect of  $\beta\gamma$  subunits. In the presence of lower concentrations of detergent,  $\text{IC}_{50}$  values of GTP-binding proteins on the enzyme activity were much less than those in the presence of 0.1% Lubrol PX.




Fig.1. Effect of GTP-binding proteins and their subunits on phosphodiesterase. Phosphodiesterase activity was measured in the reaction mixture containing 50 mM Tris-HCl (pH 7.5), 5 mM  $\text{MgCl}_2$ , 1 mM DTT, 0.1 mg/ml bovine serum albumin, 2  $\mu\text{M}$  cyclic [ $^3\text{H}$ ]AMP, 0.1% Lubrol PX, 10 nM calmodulin, 0.5 mM  $\text{CaCl}_2$  (●, ▲, ■) or 0.5 mM EGTA (○, □), and varying concentrations of GTP-binding proteins,  $G_i$  or  $G_o$  (●, ○),  $\alpha$  subunit (▲), and  $\beta\gamma$  subunits (■, □). The reaction was started by the addition of the calmodulin-deficient phosphodiesterase preparation (0.22  $\mu\text{g}$  protein), and was carried out for 10 min at 30°C.

For example,  $\text{IC}_{50}$  values of  $G_i$  and  $G_o$  were 35 and 65 nM, respectively, at a Lubrol concentration of 0.0025%.

The inhibitory effects of  $G_o$  on calmodulin-stimulated phosphodiesterase activity could be overcome by increasing concentrations of calmodulin (fig.2); higher concentration of calmodulin was required to antagonize the inhibitory effects of 4  $\mu\text{M}$   $G_o$  than of 1  $\mu\text{M}$   $G_o$ . Similar results were observed with  $G_i$  (not shown).

To determine whether GTP-binding proteins bind directly to calmodulin, the gel permeation binding experiment of Hummel and Dreyer [22]




Fig. 2. The reversal of the inhibitory effect of  $G_0$  on calmodulin-stimulated phosphodiesterase by increasing concentrations of calmodulin. Phosphodiesterase activity was measured in the reaction mixture containing 0.5 mM  $\text{CaCl}_2$  and varying concentrations of calmodulin in the presence (●, 1  $\mu\text{M}$ ; ▲, 4  $\mu\text{M}$ ) or absence (○) of  $G_0$ .

was carried out with Sephadex G-50 column equilibrated with [ $^3\text{H}$ ]calmodulin. Fig. 3 illustrates a typical chromatogram obtained in a  $\text{Ca}^{2+}$ -containing buffer following the application of  $G_0$ . Fig. 3 also shows that the binding of  $G_0$  to calmodulin appears  $\text{Ca}^{2+}$ -dependent, since EGTA prevents the formation of the  $G_0$ -calmodulin complex. Similar results were observed with  $G_i$  (not shown). These results indicate that GTP-binding proteins bind directly to calmodulin in a  $\text{Ca}^{2+}$ -dependent manner.

#### 4. DISCUSSION

Both  $G_i$  and  $G_0$  prevented the activation of cyclic nucleotide phosphodiesterase by calmodulin. The results from the gel permeation binding experiments of Hummel and Dreyer [22] indicated that GTP-binding proteins bound to calmodulin in a  $\text{Ca}^{2+}$ -dependent manner.  $G_0$  was a less potent inhibitor of phosphodiesterase than  $G_i$ , but it exists at much higher concentration in the brain membranes. Because the inhibitory effect of  $G_0$  was derived from  $\beta\gamma$  subunits,  $\beta\gamma$  subunits should be the major molecules to interact with calmodulin. The affinities of the GTP-binding proteins for calmodulin are not so high as that of calmodulin-regulated enzyme or various




Fig. 3. Elution profiles of  $G_0$  on Sephadex G-50 column equilibrated with the buffer containing [ $^3\text{H}$ ]calmodulin. Sephadex G-50 column (0.7  $\times$  11 cm) was equilibrated with the buffer containing 10 mM Hepes (pH 8.0), 0.5 mM DTT, 0.05% Lubrol PX, [ $^3\text{H}$ ]calmodulin and 0.5 mM  $\text{CaCl}_2$  or 0.5 mM EGTA at 25°C.  $G_0$  (2 nmol) was applied to the column and each fraction was analyzed for radioactivity.

cytoskeletal calmodulin-binding proteins [4]. But the  $\text{IC}_{50}$  values of  $G_i$  and  $G_0$  were lower than those of opioid peptides [11], and were the same order with insect venom peptides [13] and seminalplasmin [14] which were recently reported to interact with calmodulin.

All assays of phosphodiesterase in this study were carried out in the presence of 0.1% Lubrol PX. When the assay was performed in the presence of lower concentrations of detergent,  $\text{IC}_{50}$  values of  $G_i$  and  $G_0$  were less than those in the presence of 0.1% Lubrol. These phenomena suggest a hydrophobic interaction between calmodulin and GTP-binding proteins. Furthermore, the  $\beta\gamma$  subunits seem to be hydrophobic proteins because they aggregate in the absence of detergent, while  $G_i\alpha$  and  $G_0\alpha$  subunits remain monomeric in the

same conditions [23]. Therefore, the inhibition of calmodulin-stimulated phosphodiesterase by  $\beta\gamma$  is probably due to the hydrophobicity of this dimer like the inhibition obtained with neuroleptics [24].

$G_i$  and  $G_o$  account for 1.5% of the membrane proteins [1], viz. 15  $\mu\text{g}/\text{mg}$  of membrane proteins. On the other hand, the concentration of calmodulin in the particulate fractions of rat cerebral cortex was reported to be 2.7  $\mu\text{g}/\text{mg}$  of membrane proteins [5]. This is a similar concentration to that of GTP-binding proteins.

The physiological significance of the interaction of calmodulin with GTP-binding proteins is still unclear. However, it may suggest a novel function of calmodulin and GTP-binding proteins because both proteins exist at high concentration in the brain membranes. It presumably includes complex regulatory system related to  $\text{Ca}^{2+}$  mobility in the cell membranes.

#### ACKNOWLEDGEMENT

This work was supported in part by research grant from the Ministry of Education, Science and Culture of Japan.

#### REFERENCES

- [1] Sternweis, P.C. and Robishaw, J.D. (1984) *J. Biol. Chem.* 259, 13806–13813.
- [2] Neer, E.J., Lok, J.M. and Wolf, L.G. (1984) *J. Biol. Chem.* 259, 14222–14229.
- [3] Gilman, A.G. (1984) *Cell* 36, 577–579.
- [4] Manalan, A.S. and Klee, C.B. (1984) *Adv. Cyclic Nucleotide Res.* 18, 227–278.
- [5] Kakiuchi, S., Yasuda, S., Yamazaki, R., Teshima, Y., Kanda, K., Kakiuchi, R. and Sobue, K. (1982) *J. Biochem.* 92, 1041–1048.
- [6] Sobue, K., Ichida, S., Yoshida, H., Yamazaki, R. and Kakiuchi, S. (1979) *FEBS Lett.* 99, 199–202.
- [7] DeLorenzo, R.J. (1982) in: *Calcium and Cell Function* (Cheung, W.Y. ed.) vol.III, pp.271–309, Academic Press, New York.
- [8] Kakiuchi, S. and Yamazaki, R. (1970) *Biochem. Biophys. Res. Commun.* 41, 1104–1110.
- [9] Sano, M. and Drummond, G.I. (1981) *J. Neurochem.* 37, 558–566.
- [10] Iqbal, Z. and Ochs, S. (1980) *Ann. NY Acad. Sci.* 356, 389–390.
- [11] Sellinger-Barnette, M. and Weiss, B. (1982) *Mol. Pharmacol.* 21, 86–91.
- [12] Malencik, D.A. and Anderson, S.R. (1982) *Biochemistry* 21, 3480–3486.
- [13] Katoh, N., Raynor, R.L., Wise, B.D., Schatzman, R.C., Turner, R.S., Helfman, D.M., Fain, J.N. and Kuo, J.-F. (1982) *Biochem. J.* 202, 217–224.
- [14] Gietzen, K. and Galla, H.-J. (1985) *Biochem. J.* 230, 277–280.
- [15] Yazawa, M., Sakuma, M. and Yagi, K. (1980) *J. Biochem.* 87, 1313–1320.
- [16] Sobue, K., Muramoto, Y., Yamazaki, R. and Kakiuchi, S. (1979) *FEBS Lett.* 105, 105–109.
- [17] Sharma, R.K. and Wang, J.H. (1979) *Adv. Cyclic Nucleotide Res.* 10, 187–198.
- [18] Hidaka, H. and Asano, T. (1976) *J. Biol. Chem.* 251, 7508–7516.
- [19] Asano, T., Ui, M. and Ogasawara, N. (1985) *J. Biol. Chem.* 260, 12653–12658.
- [20] Asano, T. and Ogasawara, N. (1986) *Mol. Pharmacol.* 29, 244–249.
- [21] Northup, J.K., Smigel, M.D. and Gilman, A.G. (1982) *J. Biol. Chem.* 257, 11416–11423.
- [22] Hummel, J.P. and Dreyer, W.J. (1962) *Biochim. Biophys. Acta* 63, 530–532.
- [23] Sternweis, P.C. (1986) *J. Biol. Chem.* 261, 631–637.
- [24] Asano, M. and Hidaka, H. (1984) in: *Calcium and Cell Function* (Cheung, W.Y. ed.) vol.V, pp.123–164, Academic Press, Orlando.